

No 43.

#11

Dysentery

Mr Hill

admitted March 17. 1817

257

...the ...  
...the ...  
...the ...  
...the ...  
...the ...

...the ...  
...the ...  
...the ...  
...the ...  
...the ...

...the ...  
...the ...  
...the ...  
...the ...  
...the ...

...the ...  
...the ...  
...the ...  
...the ...  
...the ...

In co  
ed it mor  
weather;  
the histo  
avoids to b  
disease in  
of my po  
setting on  
of  
the Summ  
it had y  
In the  
from 55  
Month.  
June a  
ging, fr  
loubd H  
In Luc  
to 95, the  
August  
the same  
Sep  
early in  
About  
made its  
Among Ch

### Observations on Dysentery.

In commencing these observations, I deemed it most prudent to notice the state of the weather in the Summer of 1818. Then to give the history, and cure of the disease, afterwards to treat and explain the nature of the disease in general. Although it will be out of my power to offer any thing new or interesting on the Subject.

I believe it was generally admitted that the Summer of 1818. in Virginia, was warmer than it had for many years.

In the Month of May the Thermometer varied from 55 to 80. heavy rains fell during the Month.

June was very hot, the Thermometer ranging from 70 to 90. frequent rains with loud thunder.

In July the Thermometer varied from 75 to 95, the month clear, warm and sultry.

August was dry and hot, the Thermometer the same as in July.

September was cool, having a frost early in the month.

About the middle of May the Dysentery made its appearance, and was very fatal, among Children. as far as I could learn, the

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

disease  
Unic  
vario  
progre  
used i  
often  
of fever  
semis  
with s  
I lived  
as fever  
violence  
was ended  
by man  
cult  
stated  
and eyes  
of bile  
ad to t  
in the  
gripping  
and bu  
our and  
contin  
dark m  
they were

disease more or less prevailed throughout the United States. It frequently occurred with various degrees of violence; in many cases its progress was arrested, if proper remedies were used in the commencement. I found it very often complicated, assuming various forms of fever, particularly the intermitting and remitting fever; it was sometimes combined with Typhus. In the neighbourhood in which I lived, it was generally connected with Bilious fever, particularly in those cases that were violent; this I knew of my own knowledge, and was credibly informed of its bilious tendency by many respectable gentlemen of the faculty. Bilious appearances were not only detected in the stools, urine &c. but the skin and eyes exhibited very marked of the diffusion of bile through the system. The disease seemed to be attended with much inflammation in the lower part of the intestinal tube, severe griping, and flatulence, the calls very frequent and bilious, preceded by excrete pail. The colour and consistence of the discharges changed continually. Sometimes the stools consisted of dark mucus tinged with blood, and at times they were of a green watery colour, and having

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

a ifor  
-mion  
-nate  
-ed f  
-ients  
in the  
the m  
-owing  
part  
-occur  
-product  
-cases  
The  
-nel  
-etite  
-all, f  
-poms  
-distur  
Chill  
-ear  
was o  
more  
-ly ac  
its whe  
-ptom  
-writa



a foetid smell; indeed it is almost impossible to give a correct account of the evacuations, for in Thrush they never continued the same six hours together. In several patients, I observed nothing passed but blood in the afternoon, and bilious discharges in the morning: this perhaps might have been owing to a return of the fever in the afternoon of the day, which would naturally increase the inflammation in the bowels and produce hemorrhage from the small vessels diseased.

The general symptoms that characterized Dysentery last Summer, were loss of appetite, sickness of the stomach, vomiting &c. all, frequent pulse, foul tongue &c. these symptoms frequently came on before there was any disturbance of the bowels. In many cases, the chills became periodical, assuming the appearance of an intermittent, in others there was a periodical flush, that partook more of the remitting type. Fever more or less accompanied the disease through out its whole course, when most of the common symptoms above described became violent, great irritation seemed to pervade the whole course.

*[Faint, illegible handwriting in cursive script, likely bleed-through from the reverse side of the page.]*

of the  
of the  
appears  
all the  
new and  
consider  
longer  
few  
after the  
Mansel  
discovered  
on the  
of the  
in a Ch  
owing a  
union  
family  
a fair  
itablis  
the us  
violent  
the effe  
wound  
resting  
frame  
cause a

of the intestines, attended by great prostration  
of strength the countenance assuming a pale  
appearance, the pulse lost its force and death  
ultimately closed the scene. But when the dis-  
-eas was less violent, it was protracted to a  
considerable length, leaving the patient very  
languid, and terminating at last in bilious  
fever. This was uniformly the case with adults,  
after the disease assumed the chronic form.  
Many cases occurred in Children, where the  
disease was suddenly removed by an eruption  
on the Skin. A most remarkable instance  
of this kind happened in the neighbourhood  
in a Child of two years old; he had been labo-  
-ring under Dysentery for four weeks, his phy-  
-sician despaired of his recovery, as also did the  
family; the usual remedies in the disease had  
a fair trial, and death seemed to be the in-  
-evitable consequence; suddenly however, to  
the astonishment of all present, a most  
violent eruption of the Skin appeared; the  
effect was, that the pulse rose, became  
round and full, attended with high fever,  
respiration quick and laborious, the whole  
frame seemed enlarged, the bowels discharges  
ceased and an obstinate bilious fever followed.

This pa  
good  
use  
it was  
vers.  
sent  
was m  
Some o  
unusu  
a ha  
mat  
: us  
most s  
was the  
was m  
latter  
the, be  
feel  
to run  
in the  
be the  
many a  
thoma  
mule  
amer to

This patient recovered and I believe enjoys  
good health. I never saw or heard of the dis-  
ease affecting adults in this way, although  
it was very common among young children.  
It is worthy of remark that Dysentery  
as it appeared in Virginia last Summer,  
was marked by several new symptoms. Some  
of which have been described, but a very  
unusual symptom which I consider new was  
a pain in the shoulder resembling Rheu-  
matism, and not unfrequently a suppu-  
-ration of Urin.

I am unable to say what situations were  
most subject to this Epidemic, but believe it  
was the opinion of Medical Men, that the disease  
was most prevalent in the low land of water  
courses. Now situations seemed most unhealth-  
-thy, particularly the districts of Country sub-  
-ject to Ague and fever. As this disease seemed  
to run into the bilious or malarial and inter-  
-mitting forms of fever, may not the cause  
be the same; as far as my memory serves me  
many are of this opinion, particularly Dr  
Thomas, who wrote an essay of air, clim  
-mate and no doubt a store of much infor-  
-mation. If I may be allowed a conjecture

from a  
be of op  
normal a  
long bid  
inter,  
back i  
month, a  
tment, i  
hargate  
length of  
action, a  
and with  
more fan  
than me  
long a  
althou  
any, with  
or no fear  
even ha  
when know  
having  
disinter  
go on to  
in games  
by h  
a part:

from what I discovered last summer, I should  
be of opinion that the cause which produces  
vernal and autumnal intermitting and remitt-  
ing bilious fever, are the same, that, produces Dys-  
entery, but I acknowledge myself unable to  
prove it. The above disease was treated dis-  
crepantly by different Physicians, but the treat-  
ments which I found most successful, was  
purgatives continued at long intervals, for a  
length of time, interposing mucilaginous injec-  
tions and drinks. Calomel was used freely  
and with much success, it seemed to act  
more forcibly on the coats of the intestines  
than milder purgatives, and brought away  
large quantities of indurated feces. In full  
bilious habits, blood was taken from the  
arm, with success. In cases attended with little  
or no fever, the Compound powder of Opium  
enemata seemed to be of singular service,  
when proper evacuations had been previously given?

Having made <sup>some</sup> general observations on  
Dysentery, as it appeared in Virginia. I will  
go on to make observations on the disease  
in general.

By Dysentery, I understand an inflammation  
of a part or the whole of the intestinal tube,

the parts  
the inner  
I am sure  
cause has  
the (thin  
reason of  
length of  
ages, but  
seat of  
quitting

The  
with me  
present  
thrust  
with  
discharge  
first  
arguing, the  
same in  
and the  
meas, or  
course of  
cavance  
called  
Dependent  
presently



The permanent seat of the disease seems to be the inner coat called the Mucus Membrane. I am unable to determine whether the disease has ever pervaded the whole length of the alimentary Canal, but it appears reasonable, in violent cases, that the whole length of the bowels may suffer by its ravages, but the common and most obstinate seat of the Disease seems to be the large gut, viz. the Colon and Rectum.—

The disease every case seems to be attended with more or less fever, severe griping, frequent bloody stools, tenesmus, restlessness, thirst &c. the stools are not always mixed with blood, frequently in mild cases, the discharge, seems mucous, when the patient is first attacked he has frequent calls discharging, the contents of the lower bowels, the pain in a short time, becomes more severe and the stools more liquid, having a bilious, mucous, or bloody cast. Should the feces in the course of the disease, assume a natural appearance they are hard, sometimes round and called by Medical writers Sybala.

Dysentery, makes its appearance, more frequently in summer and autumn, than I have known

Several  
Barons o  
occur in  
men occa  
this is over  
where the  
troubles  
common  
with the  
the king a  
communion  
Biden how  
the content  
the house  
at hand  
consider  
tion, for  
that day  
received a  
case, sit  
less, such  
was, with  
the house  
frequently  
high. But

Several regular <sup>fevers</sup> formed in winter and spring, as  
Nauseous obstructions are assigned why it should  
occur in warm climates. I believe it is often  
occasioned by cold succeeding intense heat.  
this is one of the causes which is said to pro-  
duce the disease. I have reference here to cold  
suddenly applied to the body, and not a con-  
tinuance of it for any length of time, when  
cold suddenly succeeds heat, perspiration is  
checked, and there may be a particular deter-  
mination to the bowels, by occluding the disease.  
Ridemann seems to be of this opinion, for  
he contends that it is a fever, thrown upon  
the bowels in consequence of obstructed  
and perspiration. A long drought may be  
considered as an other cause of its produc-  
tion, for we find it prevailing sometimes in  
hot dry weather. Moisture, this may be con-  
sidered as an other exciting cause of the dis-  
ease, situations much exposed to heavy  
dews, such as the neighbourhood of large riv-  
ers, mill ponds &c. are all liable to obstruc-  
tion of the bowels. Complaints. Bad water very  
frequently produces the complaint if used  
freely. Rotten or unhealthy food. This is also

Said to  
uses the  
which  
and, Sam  
the  
this was  
dity. N  
men of  
men  
written  
sphere  
me app  
unrec  
entire  
row of  
Depo  
be can  
many  
areas  
its  
with  
seems  
as, the  
to take  
would  
ling, be

Said to have its influence, and no doubt, pro-  
duces the disease, in many cases; But the cause  
which most frequently gives rise to Dysente-  
ry, seems to be specific contagion, and when  
a neighbourhood once becomes affected in  
this way, it spreads with astonishing rapidi-  
ty. Not unfrequently the disease up, seems  
as an Epidemic; owing, I suppose, to a pec-  
uliar condition of the Atmosphere. It seems  
evident in many instances that the Atmo-  
sphere is the medium through which we  
are assailed by many illness. I think it not  
unreasonable to suppose that it may be pos-  
sible to find certain principles favourable to the produc-  
tion of Dysentery.

Dysentery seems to be ~~considered~~ a high-  
ly contagious disease, it is acknowledged by  
many Medical Men, to be infectious, when the  
disease appears in a family, it generally affe-  
cts many of its members, some slightly, others  
with severity, This circumstance with others  
seems almost conclusive, that it is infectio-  
us. This being admitted, great care should  
be taken to arrest the exciting cause. It  
would be well therefore to change the sit-  
ing, frequently, as also the patient's linen;

to move  
the house,  
these days  
these means  
of even less  
indication  
on in general  
acts. If the  
with much  
judge it not  
however and  
cause are very  
less of the law  
but I am in  
have come in  
has been of S  
sides as low  
symptoms se  
up towards  
of panga live  
kept open lo  
time, and  
most of the  
this course I  
such as warm

to remove the discharges some distance from the house, fumigate the room two or three times daily, with turpentine vinegar &c. &c. these means some of the exciting causes of dysentery may be removed.

As to respects the cure of dysentery, the first indication seems to be, to lessen the inflammation in general, by bleeding and other evacuations. If the patient should be plethoric, with much fever, full rounded pulse, I should judge it necessary to bleed frequently. I am however aware that many writers, on this disease are very cautious with respect to the use of the lancet, and perhaps with good reason, but I am well assured from the few cases that have come under my observation, that bleeding has been of singular service. It should be repeated as long as the pulse and other febrile symptoms seem to require. The second step towards a cure, seems to be, the exhibition of purgative remedies. The bowels should be kept open freely with mild purgatives, continued daily for a length of time, viz until most of the febrile symptoms are removed. This course should be assisted with injections, such as warm milk, and gruel mucilage of Gum

le me  
- cul  
be ne  
if we  
- m  
Houl  
als m  
Hytte  
so be  
it seen  
much  
If  
some  
ation  
tion  
of grace  
have  
ma  
the  
some  
Small  
some  
in the  
some  
of spec



drastic, sherry<sup>and</sup> &c. the patients drink should be mild and at the same time nourishing and the-  
cilaginous. After proper evacuations, it will be well to determine to the surface of the body, by using rubefacient diaphoretic remedies, for which  
bainrose, small doses of Antimonial Wine may  
should be used and continued at proper inter-  
vals, in order to keep up a gentle nausea and  
slight perspiration. Specuacum has most al-  
so be of singular service in the cure, for  
it seems to be a medicine that acts with  
much force on the skin.

If the Abdomen be tumid, hard and  
sore to the touch, in addition to fomenta-  
tions and anodyne liniments, the applica-  
tion of a blister to it, will be found to be  
of great service, when the febrile symptoms  
have been reduced, and the bowels irregularly  
evacuated, Opium must be administered,  
for the purpose of relieving pain, and in-  
creasing sleep. Luleanum combined with  
small doses of Antimonial wine, will afford  
and much relief. Opium and Specuacum  
in the form of pill or powder, has also been given  
in with advantage. The Compound Powder  
of Specuacum, is an excellent remedy. Opium

From C  
and me  
of March  
the king  
of the pr  
the  
large the  
action  
on land  
out of the  
mishab.  
Tom  
entices  
ation of  
ite or  
in admi  
of Sy  
to be a  
shall be  
Shirato  
long is  
The  
aimen C  
onance  
the Shin  
fish. of

Should be administered at long intervals, and not then unless the patient complains of much pain or restlessness and want of sleep. This however should depend upon the judgment of the practitioner.

I have made no mention of Emetics, although they are spoken of by some writers, under certain circumstances they may be of singular service, perhaps in cases where there is great disorder of the stomach, they may be useful.

Tonics and astringents may be used in cases of extreme weakness. The infusion of Cinchona, Columbo, Chamomile, or Gentian with Aromatics, may be administered freely in ~~the~~ <sup>the</sup> ~~asthenia~~ <sup>asthenia</sup> of Dysentery. Whenever there appears to be any acidity in the stomach, the Chalk Lulap, or any mild absorbent, should be ordered and continued as long as necessity requires.

The Article of Cloathing, in the treatment of Dysentery, is of no small importance. Flannel should be worn next the skin in general, with one or two folds of it over the Abdomen. This answers

*[Faint, mostly illegible handwriting in cursive script, covering the majority of the page. The text appears to be a continuous narrative or letter.]*

the two  
tion, and  
the weak  
interic  
and easy  
Substan  
howley a  
hic dispo  
Shipping

11

the twofold purpose of promoting perspiration, and giving the necessary support to the weakened intestines. The diet of Dysenteric patients, should be very light and easy of digestion. Mucilaginous substances should be freely given, as barley water, Rice water, Gum Arabic dissolved in water, Arrow root, Slippery elm tea Jellies &c.

The first thing I saw when I  
came out of the house was  
the morning sun. It was so  
bright that I had to close my eyes  
for a moment. I was so happy  
to see the sun. I had been  
in the house for so long  
that I had almost forgotten  
what it was like. I was so  
glad to see the sun. I had  
been so sad. I had been so  
lonely. I had been so alone.  
I had been so happy. I had  
been so sad. I had been so  
lonely. I had been so alone.

100  
00  
The con